

Overactive Bladder

Overactive Bladder (OAB) is a complex condition affecting about 15% of women across all ages. It can be costly, embarrassing and result in avoiding social activities for fear of leakage. Age increases the risk for OAB as do some bladder conditions (infection, bladder stones, or abnormal growths). For some women, the cause is unknown.

The Overactive Bladder


Women with OAB feel a sudden urge to urinate, sometimes followed by leaking. Some women leak on the way to the bathroom, or while they are taking down their clothes. Sometimes the whole bladder empties. It is common to feel the frequent urge to go even though your bladder isn't full.

Your bladder is a sack of muscle that holds urine. As urine flows into the bladder, the walls expand to make room for more fluid, like a water balloon. The muscles that surround your urethra, the tube you urinate from, tighten to hold in urine while the bladder muscle relaxes.


When you urinate, your brain signals the urethra and pelvic floor muscles to relax. The bladder muscle squeezes, pushing the urine out of the body. Typically, women urinate four to seven times per day and one time at night.

In women with OAB, the bladder muscle is overactive and squeezes too often. These bladder spasms make it feel like you need to urinate often and quickly, even if there isn't much urine in your bladder.

NORMAL BLADDER



OVERACTIVE BLADDER



The muscle that lines the bladder, called the detrusor muscle, squeezes down too often and too powerfully. This causes urine to come out at the wrong times. Common things that bring this on are hearing running water, putting your key in the door at home, or seeing the toilet.

©Allia Medical Media

LEARN THE TERMS

Urinary urgency (“gotta go right now”): Sudden, strong desire to pass urine, which is very difficult to defer.

Urinary frequency (“gotta go often”): Urinating eight or more times per day.

Nocturia: Need to urinate one or more times during sleeping hours.

Urinary urgency incontinence (UUI): Urinary leakage that occurs with the sudden, strong desire to pass urine.

Overactive bladder: Urinary urgency, usually with frequency and nocturia, and sometimes with urgency urinary incontinence. This occurs without an infection or other health problem.

Diagnosis

It can feel embarrassing to have these bladder issues. You are not alone. Talk to your urogynecologist (urogyn) as a first step to taking back your life. Your doctor will ask questions about when and how often you leak urine. A physical exam helps the doctor identify other conditions that influence the bladder, such as prolapse.

Additional tests might include:

- Urine analysis to check for a urinary tract infection and blood in your urine.
- Ultrasound to assess how much urine remains in your bladder after urinating.
- Urodynamics to provide information on your bladder and urethra.

Overactive Bladder

The doctor may ask you to complete a bladder diary. This requires you to record what, how much, and how often you drink. You also measure the amount you urinate. This will help your doctor learn more about your bladder symptoms.

Treatments

Ask your urogyn about the best treatment or combination of therapies for you. OAB can be controlled, and you can get your life back.

DIETARY CHANGES

Certain types of drinks can provoke bladder spasms. For example, coffee, tea, sodas and other beverages with caffeine worsen symptoms. Artificial sweeteners, fruit juices and alcohol can also bother your bladder. Completing a bladder diary can help you look for a connection between your diet and bladder symptoms.

PADS AND SKIN CARE

Hopefully OAB treatment will reduce leakage. In the meantime, use only all-cotton or incontinence pads (not menstrual pads), which are gentle on the skin. Try to let the skin air out and dry when possible. Some women need to use a barrier cream to protect the skin near the vagina.

BLADDER TRAINING

Women with OAB tend to go to the bathroom often. However, this can make symptoms worse. Your bladder learns to hold less urine, leading to even more frequent bathroom trips. Bladder re-training involves using your pelvic floor muscles and “mind over bladder” techniques to gradually increase time between bathroom visits. As you do this more and more, your bladder muscle readjusts. This allows you to make fewer bathroom trips.

This retraining can be hard to do – ask your doctor for a referral to a pelvic floor physical therapist (PT) to learn how to do it.

MEDICINES

OAB medicines help your bladder hold more urine for longer periods of time. They also reduce urine leakage. Potential side effects include dry mouth and constipation. There are many different medicines available. Let your doctor know how the drugs work for you. It

may take trying different medicines before you find the one that works best for you.

PELVIC FLOOR PHYSICAL THERAPY

Most women find that pelvic floor muscle exercises help improve symptoms. For the best effect, work with a specialized PT to learn the techniques. Ask your doctor for a referral. Then use these exercises every time you feel urgency. It may take 3 to 6 months of regular pelvic floor muscle exercise to see results.

BOTULINUM TOXIN (BOTOX)

Botox relaxes the bladder muscle. This allows more urine to be held in the bladder before you have to go to the bathroom. Under a local anesthetic or sedation, your doctor uses a small camera and needle to inject Botox into the bladder wall.

Typically, the injection needs to be repeated 1-2 times a year. A small number of women experience side effects, such as difficulty emptying their bladder or urinary tract infection. Ask your doctor if Botox is right for you.

NERVE STIMULATION

There are many nerves involved in bladder function. Nerve stimulators help control these nerves, reducing the need to urinate often.

Tibial Nerve Stimulation is an external device. A small needle is inserted near a nerve in the ankle during a series of office visits. When connected to the device, nerve messages are altered.

Sacral Nerve Stimulation is a device implanted near the spine. It modulates the bladder nerves.

Three Takeaways

- 1. Bladder spasms can create the need to urinate urgently and often.**
- 2. Diet changes and bladder retraining can reduce OAB. A bladder diary can help provide more information about the link between diet and your bladder symptoms.**
- 3. Treatments, including physical therapy, medications, Botox, and nerve stimulation, help manage the symptoms.**